


## *Graphidium strigosum*

*Esther van Praag, Ph.D.*

This cosmopolitan parasite is mainly observed in the wild rabbit (*Oryctolagus cuniculus*), and in palaeartic Leporidae, such as the hare (*Lepus europaeus*, *L. capensis*). The rabbit is probably the original host, as it tolerates the presence of *G. strigosum* much better than the hare, which presents important stomach lesions. It is thus speculated that the presence of this worm in hare populations depends on the presence of rabbits. The relationship of *Graphidium strigosum* and *Trichostrongylus retortaeformis* has been reported for wild rabbits. House rabbits become infested by green forage, contaminated by infective larvae.

Not much is known about the biology and the life cycle of this parasite.



The eggs are approximately 95\*50 µm in size and are excreted in the morula stage. Under good environmental conditions, the larvae hatch about 10 h. later. The L<sub>2</sub> stage is reached in a further 2-3 days. Ensheathed L<sub>3</sub> larvae are infective and migrate along the herbage according to the period of the day: to the tip at dusk, downwards when exposed to sun light and heat, till they are ingested by their host. Male and female adult worms are red with many longitudinal lines and transversal striations. The males measure about 12 mm long, while females average 16 mm. The male possess paired and slender spicules and a well-developed copula-bursa.

### Symptoms

Clinical signs are close to those of gastritis. Massive infestations can cause catarrhal gastritis with some fibrosis, and extreme inflammation of various parts of the intestinal tract (stomach, small intestine, cecum).

Necropsy shows that L<sub>4</sub> stage worms are coiled in the ducts of the gastric



glands of the fundus. Adults worms are generally located in the mucus layers, with the head buried in the stomach grooves, without attachment to the mucosa.

### Treatment

<b>Benzimidazoles</b>	fenbendazole	20 mg/kg, PO, repeated after 10-14 days
<b>Macrolides</b>	ivermectin	0.4 mg/kg, SC, repeated after 10-14 days

### Further information

1. B. Boag (1987) The Helminth Parasites of the Wild Rabbit *Oryctolagus cuniculus* and the brown hare *Lepus capensis* from the Island of Coll, Scotland. J. Zool. 212: 352-355.
2. B. Boag and H.H. Kolb (1989) Influence of the Host Age and Sex on Nematode Populations in the Wild Rabbit (*Oryctolagus cuniculus* L.). Proc. Helminth. Soc. Washington 56: 116-119.
3. Brookhuizen and Kemmers (1976) The Stomach Worm *Graphidium strigosum* (Dujardin) Railliet and Henry, in the European Hare, *Lepus europaeus* Pallas. In: Pielowski, Z. and Pucek Z. (eds) Ecology and Management of the European Hare Populations. Panstwowe Wydawnictwo Rolnicze i Lesne, Warsaw, Poland, pp 157-171.
4. J.D. Dunsmore, M.L. Dudzinski (1968) Relationship of Numbers of Nematode Parasites in Wild Rabbits, *Oryctolagus cuniculus* (L.), to Host Sex, Age and Season. J. Parasitol. 54: 462-474.
5. E.A. Nickel and W. Haupt (1986) Experimental Studies on the Course and Consequences of Infection with *Graphidium strigosum* (Nematoda, Trichostrongylidae) in *Oryctolagus cuniculus*. Agnew. Parasitol. 27, 215-219.
6. E.J.L. Soulsby (1968) "Helminths, Arthropods, and Prorozoa of Domestic Animals". Williams and Wilkins, Baltimore, Maryland.
7. R. Wetzel, K. Enigk (1937) Zur Biologie von *Graphidium strigosum*, dem Magenwurm der Hasen und Kaninchen. Deutsch. Tierärztliche Wochenschrift 45: 401-405.

---

*The information on these pages may not be reproduced, or republished on another webpage, website, or elsewhere.*

FEBRUARY 2003

